Opposite Action Readiness - Clinician dialogue

Guide a collaborative conversation about opposite action readiness while checking fit and participant choice.

Mental health professionals | 15 minutes

Fictional example: Taylor wants to avoid a harmless routine question after assuming annoyance.

Invite and obtain permission

Offer: "Would it be useful to explore opposite action readiness using an ordinary or fictional example? We can adapt it, observe, or stop." Ask what the person wants from the conversation.

Elicit before teaching

Ask: "How does this show up, if at all, in your own words?" Offer this idea tentatively: Changing an emotional action urge depends on fit, facts, and safety. Ask what fits and what does not.

Guide the task

With a clinician, assess whether an urge fits the facts and whether acting on it is effective before designing a small alternative. Ask for permission before giving an example. Use this fictional illustration if useful: Taylor wants to avoid a harmless routine question after assuming annoyance.

Reflect and check accuracy

Reflect only what was offered: "I heard you describe ___. What did I miss?" Ask: What makes an alternative action appropriate in this specific situation? Avoid assigning motives or interpreting silence as agreement.

Respond to a mismatch

If the person rejects the frame, say: "Thank you for correcting it. What would be more useful?" Return to their language or choose a different approach.

Close with agency

Ask which observation, question, or support request the person wants to keep. Confirm a manageable ending and appropriate follow-up; do not turn a tentative possibility into a homework obligation.

Reflection

What would tell me to adapt, stop, or seek supervision?

Scope and adaptation

These are selected DBT-informed educational exercises, not comprehensive DBT, phone coaching, or crisis care. Use individualized clinical assessment and existing care plans. No cold-water immersion, breath holding, intense exercise prescriptions, exposure assignments, or risk scoring are included. Behavior chains and opposite-action decisions need clinician guidance when risk or complexity is present.

Original AI-assisted educational draft; independent clinical review pending. Version 1.0. Background: Dialectical Behavior Therapy Skills Training: Fundamentals. Full source links and scope: library.morethantherapy.org/approaches/resources/DBT-A13-3