Experiment by Consent - Session activity

Facilitate a structured practice of experiment by consent with preparation, reflection, and an agreed closing.

Mental health professionals | 30 minutes

Fictional example: Rae tries asking a question more slowly in a fictional conversation.

0-5 min | Agree on purpose and fit

Explain the educational scope. Ask what would be useful, check readiness and access needs, and establish choices to observe, adapt, or pass. Use an ordinary example; do not solicit trauma narratives.

5-10 min | Orient with an example

An experiment is an invitation to learn rather than a performance test. Fictional example: Rae tries asking a question more slowly in a fictional conversation. Ask what is observed and what remains uncertain.

10-20 min | Practice and compare

Design a reversible 30-second change in wording, seating preference, or pace and agree what to notice. Offer quiet individual work, a supported conversation, or an observing role. Compare two possible responses without evaluating people as right or wrong.

20-25 min | Reflect on meaning

What did you learn, including reasons not to repeat it? Ask what fit, what did not fit, and how culture, power, disability access, or external conditions affect the options.

25-30 min | Integrate and close

Invite one optional next step or support request. Orient to the current setting, allow an unfinished question to remain unfinished, and agree how to revisit it. End early or adapt if activation increases.

Adaptations

For a group, use fiction and no compulsory round-robin disclosure. For remote use, allow off-camera reflection and private notes. For brief use, keep one example and one question while retaining opening and closing choices.

Reflection

What adaptation or follow-up is needed?

Scope and adaptation

Keep experiments collaborative and low intensity. Chair-based or unfinished-business exploration belongs with appropriately trained clinicians after assessment, with a stop cue and an observing alternative. Do not interpret symbols, recover memories, force body focus, or use cathartic confrontation.

Original AI-assisted educational draft; independent clinical review pending. Version 1.0. Background: Gestalt Therapy: An Introduction. Full source links and scope: library.morethantherapy.org/approaches/resources/GST-A12-4